



A PATIENT'S GUIDE TO CARPAL TUNNEL SYNDROME



INTRODUCTION

Carpal tunnel syndrome (CTS) is a common problem affecting the hand and wrist. Symptoms begin when the median nerve gets squeezed inside the carpal tunnel of the wrist, a medical condition known as nerve entrapment. Any condition that decreases the size of the carpal tunnel or enlarges the tissues inside the tunnel can produce the symptoms of CTS. This syndrome has received a lot of attention in recent years because of suggestions that it may be linked with occupations that require repeated use of the hands, such as typing on a computer keyboard or doing assembly work. Actually, many people develop this condition regardless of the type of work they do.

This guide will help you understand:

- where the carpal tunnel is located
- how CTS develops
- what can be done for the condition

ANATOMY

Where is the carpal tunnel, and what does it do?

The carpal tunnel is an opening through the wrist to the hand that is formed by the bones of the wrist on one side and the transverse carpal ligament on the other. (Ligaments connect bones together.) This opening forms the carpal tunnel.

The median nerve passes through the carpal tunnel into the hand. It gives sensation to the thumb, index finger, long finger, and half of the ring finger.



It also sends a nerve branch to control the thenar muscles of the thumb. The thenar muscles help move the thumb and let you touch the pad of the thumb to the tips each of each finger on the same hand, a motion called opposition. The median nerve and flexor tendons pass



NEXT PAGE ►

ABOUT MR GEORGE-MALAL


Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS


SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com



through the carpal tunnel. The median nerve rests on top of the tendons, just below the transverse carpal ligament. The flexor tendons are important because they allow movement of the fingers, thumb, and hand, such as when grasping. The tendons are covered by a material called tenosynovium. The tenosynovium is a slippery covering that allows the tendons to glide next to each other as they are worked.



CAUSES

What causes CTS?

Any condition that makes the area inside the carpal tunnel smaller or increases the size of the tissues within the tunnel can lead to symptoms of CTS. For example, a traumatic wrist injury may cause

swelling and extra pressure within the carpal tunnel. The area inside the tunnel can also be reduced after a wrist fracture or dislocation if the bone pushes into the tunnel.

Any condition that causes abnormal pressure in the tunnel can produce symptoms of CTS. Various types of arthritis can cause swelling and pressure in the carpal tunnel. Fractured wrist bones may later cause CTS if the healed fragments result in abnormal irritation on the flexor tendons. Other conditions in the body can produce symptoms. Pregnancy can cause fluid to be retained, leading to extra pressure in the carpal tunnel. Diabetics may report symptoms of CTS, which may be from a problem in the nerve (called neuropathy) or from actual pressure on the median nerve. People with low thyroid function (called hypothyroidism) are more prone to problems of CTS.

The way people do their tasks can put them at more risk for problems of CTS. Some of these risks include:

- force
- posture
- wrist alignment
- repetition
- temperature
- vibration

One of these risks alone may not cause a problem. But doing a task that involves several factors may pose a greater risk. And the longer a person is exposed to one or more risks, the greater the possibility of having a problem with CTS. However, scientists believe that other factors such as smoking, obesity, and caffeine intake may actually be more important in determining whether a person is more likely to develop CTS. In other instances, CTS can start when the tenosynovium thickens from irritation or inflammation. This thickening causes pressure to build inside the carpal tunnel.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL


Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com



But the tunnel can't stretch any larger in response to the added swelling, so the median nerve starts to squeeze against the transverse carpal ligament. If the pressure continues to build up, the nerve is eventually unable to function normally.

When pressure builds on the median nerve, the blood supply to the outer covering of the nerve slows down and may even be cut off.


The medical term for this is ischemia. At first, only the outside covering of the nerve is affected. But if the pressure keeps building up, the inside of the nerve will start to become thickened. New cells (called fibroblasts) form within the nerve and create scar tissue. This is thought to produce the feelings of pain and numbness in the hand. If pressure is taken off right away, the symptoms will go away quickly. Pressure that isn't eased right away can slow or even stop the chances for recovery.

SYMPTOMS

What does CTS feel like?

One of the first symptoms of CTS is gradual tingling and numbness in the areas supplied by the median nerve. This is typically followed by dull, vague pain where the nerve gives sensation in the hand. The hand may begin to feel like it's asleep, especially in the early morning hours after a night's rest.

Sometimes pain may even spread up the arm to the shoulder. If the condition progresses, the thenar muscles of the thumb can weaken, causing the hand to be clumsy when picking up a glass or cup. If the pressure keeps building in the carpal tunnel, the thenar muscles may begin to shrink (atrophy).



Touching the pad of the thumb to the tips of the other fingers becomes difficult, making it hard to grasp items such as a steering wheel, newspaper, or telephone.

DIAGNOSIS

How do doctors identify the condition?

Your doctor begins the evaluation by obtaining a history of the problem, followed by a thorough physical examination. Your description of the symptoms and the physical examination

NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT


EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

are the most important parts in the diagnosis of CTS. Commonly, patients will complain first of waking in the middle of the night with pain and a feeling that the whole hand is asleep.

Careful investigation usually shows that the little finger is unaffected. This can be a key piece of information to make the diagnosis. If you awaken with your hand asleep, pinch your little finger to see if it is numb also, and be sure to tell your doctor if it is or isn't. Other complaints include numbness while using the hand for gripping activities, such as sweeping, hammering, or driving.

If your symptoms started after a traumatic wrist injury, X-rays may be needed to check for a fractured bone.

If more information is needed to make the diagnosis, electrical studies of the nerves in the wrist may be requested by your doctor. Several tests are available to see how well the median nerve is functioning, including the nerve conduction velocity (NCV) test. This test measures how fast nerve impulses move through the nerve.



TREATMENT


What can be done for CTS?

Non-Surgical Treatment

Activities that are causing your symptoms need to be changed or stopped if at all possible.

Avoid repetitive hand motions, heavy grasping, holding onto vibrating tools, and positioning or working with your wrist bent down and out. If you smoke, talk to your doctor about ways to help you quit. Lose weight if you are overweight. Reduce your caffeine intake.

A wrist brace will sometimes decrease the symptoms in the early stages of CTS. A brace keeps the wrist in a resting position, not bent back or bent



down too far. When the wrist is in this position, the carpal tunnel is as big as it can be, so the nerve has as much room as possible inside the carpal tunnel. A brace can be especially helpful for easing the numbness and pain felt at night because it can keep your hand from curling under as you sleep. The wrist brace can also be worn during the day to calm symptoms and rest the tissues in the carpal tunnel.

Anti-inflammatory medications may also help control the swelling and reduce symptoms of CTS. These include common over-the-counter medications such as ibuprofen and aspirin. Oral steroid medication may also offer some relief. In some studies, high doses of vitamin B-6 have been shown to help in decreasing CTS symptoms.

Some types of exercises have also shown to help prevent or at least control the symptoms of CTS. If these simple measures fail to control your symptoms, an injection of cortisone into the carpal tunnel may be suggested. This medication is used to reduce the swelling in the tunnel and may give temporary relief of symptoms.

A cortisone injection may help ease symptoms and can aid your doctor in making a diagnosis.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL


Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com



If you don't get even temporary relief from the injection, it could indicate that some other problem is causing your symptoms. When your symptoms do go away after the injection, it's likely they are coming from a problem within the carpal tunnel. Some doctors feel this is a signal that a surgical release of the transverse carpal ligament would have a positive result.

Your doctor may suggest that you work with a physical or occupational therapist. The main focus of treatment is to reduce or eliminate the cause of pressure in the carpal tunnel. Your therapist may check your workstation and the way you do your work tasks. Suggestions may be given about the use of healthy body alignment and wrist positions, helpful exercises, and tips on how to prevent future problems. You might also receive treatments to reduce inflammation and to encourage normal gliding of the tendons and median nerve within the carpal tunnel.

Surgery


If all attempts to control your symptoms fail, surgery may be suggested to reduce the pressure on the median nerve. Several different surgical procedures have been designed to relieve pressure on the median nerve. By releasing the pressure on the nerve, the blood supply to the nerve improves, and most people get relief of their symptoms.

However, if the nerve pressure has been going on a long time, the median nerve may have thickened and scarred to the point that recovery after surgery is much slower.


Open Release

The standard surgery for CTS is called open release. Open surgical procedures use a large skin incision. In open release for CTS, a sizeable incision is made down the front of the wrist and palm, usually about two inches long. By creating a large incision, the surgeon is able to clearly see the wrist structures and to carefully do the operation.

The surgeon cuts the transverse carpal ligament in order to take pressure off the median nerve.



After dividing the transverse carpal ligament, the surgeon stitches just the skin together and leaves the loose ends of the transverse carpal ligament separated. The loose ends are left apart to keep pressure off the median nerve. Eventually, the gap between the two ends of the ligament fills in with scar tissue.



NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD


CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

Endoscopic Release

Some surgeons are using a newer procedure called endoscopic carpal tunnel release. The surgeon merely nicks the skin in order to make one or two small openings for inserting the endoscope. An endoscope is a thin, fibre-optic TV camera that allows the surgeon to see inside the carpal tunnel as the transverse carpal ligament is carefully released.

Upon inserting the endoscope, the surgeon can see the wrist structures on a TV screen. A special knife is used to cut only the transverse carpal ligament. The palmar fascia and the skin over the wrist are not disturbed.



As in open release, the loose ends of the transverse carpal ligament are left apart after endoscopic release to keep pressure off the median nerve. The gap eventually fills in with scar tissue.

REHABILITATION

What should I expect after treatment?

Non-Surgical Rehabilitation

If nonsurgical treatment is successful, you may see improvement in four to six weeks. You may need to continue wearing your wrist splint at night to

control symptoms and keep your wrist from curling under as you sleep. Try to do your activities using healthy body and wrist alignment. Limit activities that require repeated motions, heavy grasping, and vibration in the hand.

AFTER SURGERY

It generally takes longer to recover after open carpal tunnel release. Pain and symptoms usually begin to improve, but you may have tenderness in the area of the incision for several months after surgery.

When the stitches are removed, your surgeon may have you work with a physical or occupational therapist for six to eight weeks.

Treatments are used at first to ease pain and inflammation. Gentle massage to the incision can help reduce sensitivity in and around the incision and limit scar tissue from building up. Special exercises are used to encourage normal gliding of the tendons and median nerve within the carpal tunnel.

As you progress, your therapist will give you exercises to help strengthen and stabilise the muscles and joints in the hand, wrist, and arm.

Other exercises are used to improve fine motor control and dexterity of the hand. Your therapist will work with you to help you do your daily and work activities safely and with the least amount of strain on your wrist and hand.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

NOTES:

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com