

A PATIENT'S GUIDE TO OSTEOARTHRITIS OF THE ELBOW



INTRODUCTION

The elbow joint is injured less often than many other joints in the body. The most common injuries of the elbow joint are fractures and dislocations. Most elbow injuries tend to heal pretty well. However, an elbow injury can lead to problems later in life. The injury changes the way the joint works just enough to cause extra wear and tear to the surfaces of the joint. Over time, the joint degenerates, causing pain and difficulty with daily activities.

This condition is called osteoarthritis, degenerative arthritis, or post-traumatic arthritis.

This guide will help you understand:


- how elbow osteoarthritis develops
- how elbow osteoarthritis is diagnosed
- what can be done to ease the pain and regain elbow movement

ANATOMY


What parts of the elbow are affected?

The elbow joint is made up of three bones: the humerus bone of the upper arm, and the ulna and radius bones of the forearm.


The ulna and the humerus meet at the elbow and form a hinge. This hinge allows the arm to straighten and bend. The large triceps muscle in the back of the



arm attaches to the point of the ulna (the olecranon). When the triceps muscle contracts, it straightens out the elbow. The biceps muscles in the front of the arm contract to bend the elbow.



The connection of the radius to the humerus allows the forearm to rotate. The upper end of the radius is round. It turns against the ulna and the humerus as the forearm and hand turn from palm down (pronation) to palm up (supination).



NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT


EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

In the elbow joint, the ends of the bones are covered with articular cartilage. Articular cartilage is a slick, smooth material. It protects the bone ends from friction when they rub together as the elbow moves. Articular cartilage is soft enough to act as a shock absorber. It is also tough enough to last a lifetime, if it is not injured.

CAUSES

What causes osteoarthritis?

Osteoarthritis is caused by degeneration of the articular cartilage of a joint. Degeneration is wear that happens over time. Doctors use the term degenerative arthritis to describe the wear and tear of a joint over many years. Degenerative arthritis is another term for osteoarthritis.



Some doctors use the term degenerative arthrosis. (Arthrosis just means that the joint is wearing out.) Arthritis is technically a condition of joint inflammation. Often, joints with osteoarthritis aren't inflamed. The term arthritis should technically be used to describe only the true inflammatory conditions, such as gout, infection, and rheumatoid arthritis.

A bad sprain or fracture can actually damage the articular cartilage. The cartilage can also be bruised when too much pressure is put on the cartilage surface. The cartilage surface may not look any different. The injury often doesn't show up until months later.

Sometimes the damage to the cartilage is severe. Pieces of the cartilage can actually be ripped away from the bone. These pieces do not grow



back. Usually they must be surgically removed. If the pieces aren't removed, they may float around in the joint, causing it to catch. They can also cause a lot of pain and do more damage to the joint surfaces.

Your body does not do a good job of repairing these holes in the cartilage surface. The holes fill up with scar tissue. Scar tissue is not as slick or rubbery as the articular cartilage. An injury doesn't have to damage the cartilage to start the process of osteoarthritis.

Any injury to the elbow joint can change the way the joint works. For example, after an elbow fracture the bone fragments may not line up exactly. They heal slightly differently from their condition before the injury. Even this slight difference can cause the joint to begin the cycle of wear and tear.

A dislocation can also cause lasting damage. After the ligaments have been injured in a dislocation, the elbow joint may move differently. This change in movement alters the forces on the articular cartilage. It's just like a machine: if the mechanism is out of balance, it wears out faster.

Over many years, this imbalance in joint mechanics can damage the articular cartilage. Since articular cartilage cannot heal itself very well, the damage adds up. Finally, the joint can no longer compensate for the damage, and the elbow begins to hurt.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

SYMPTOMS

What symptoms does osteoarthritis cause?

Pain is the main symptom of osteoarthritis of any joint. At first, the pain comes only with activity. Most of the time the pain lessens while doing the activity, but after resting for several minutes pain and stiffness increase. As the condition worsens, you may feel pain even when resting. The pain may interfere with sleep.

You may have swelling around your elbow. Your elbow joint may fill with fluid and feel tight, especially after using it. When all the articular cartilage is worn off the joint surface, you may begin hearing a squeak and feel a creak in the joint when you move your elbow.

This creaking sensation is called crepitus. Osteoarthritis eventually affects the elbow's motion. The elbow joint is one of the most sensitive to injury. It quickly becomes stiff and loses motion. The first thing most people notice is that it becomes hard to completely straighten the arm. Later they find it hard to bend.

DIAGNOSIS

How will my doctor know if I have osteoarthritis?

The diagnosis of osteoarthritis of the elbow begins with a medical history of the problem. Your doctor will ask many questions about your pain, how it affects your daily activities, and any past injuries to your elbow. Because osteoarthritis develops over a long time, you may be asked questions about things you did long ago. For example, throwing athletes (such as baseball pitchers) have a higher risk of developing osteoarthritis of the elbow later in life.

Your doctor will also do a physical examination of the elbow and possibly other joints in the body. The moving and prodding may hurt, but it is important that your doctor sees exactly where and when you feel pain. Your doctor will feel for catching sensations as the joint moves.

Catching may be caused by loose fragments of cartilage and bone. (These fragments are sometimes called joint mice.)



You will probably need to get regular X-rays. X-rays are usually the best way to see what is happening with your bones. X-rays can help your doctor assess the damage and track how your joint changes over time. X-rays can also help your doctor estimate how much articular cartilage is left.

Your doctor may order blood tests if there is any question about the cause of your osteoarthritis. Blood tests can show certain systemic diseases, such as rheumatoid arthritis.

TREATMENT

What can be done to get rid of my pain?

Non-Surgical Treatment

In almost all cases, doctors try non-surgical treatments first. Surgery is usually not considered until it has become impossible to control your symptoms.

The goal of non-surgical treatment is to help you manage your pain and use your elbow without causing more harm. Your doctor may recommend non-steroidal anti-inflammatory drugs (NSAIDs), such as aspirin and ibuprofen, to help control swelling and pain. Other treatments, such as heat, may also be used to control your pain.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD


CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

Rehabilitation services, such as physical and occupational therapy, have a critical role in the treatment plan for elbow osteoarthritis. The main goal of therapy is to help you learn how to control symptoms and maximise the health of your elbow. You'll learn ways to calm your pain and symptoms. You may use rest, heat, or topical rubs.

You may be issued a special elbow splint to immobilise and protect the elbow. Resting the joint can help ease pain and inflammation. Range-of-motion and stretching exercises can improve your elbow flexibility. Strengthening exercises for the arm help steady the elbow and protect the joint from shock and stress.

Your therapist will give you tips on how to get your tasks done with less strain on the joint. To get rid of your pain, you may also need to limit your activities. You may even need to change jobs, if your work requires heavy, repetitive motions with the hand and wrist.




An injection of cortisone into the elbow joint can give temporary relief. Cortisone is a powerful anti-inflammatory medication. It can very effectively relieve pain and swelling. Its effects are temporary, usually lasting several weeks to months. There is a small risk of infection with any injection into the joint, and cortisone injections are no exception.

Surgery

Eventually, it may be necessary to consider some type of surgical treatment. There are several operations to treat advanced osteoarthritis of the elbow. Your surgeon will consider many factors when deciding which procedure is best for you, including the severity of joint degeneration, your age, your activity level, and how you use your elbow.

Arthroscopic Debridement

If you are in an early stage of osteoarthritis, your doctor may recommend arthroscopic debridement.



Arthroscopic procedures use an arthroscope, a tiny TV camera that is inserted into the joint through a very small incision. The arthroscope allows the surgeon to see inside the elbow joint.

In arthroscopic debridement, the surgeon makes other small incisions for inserting special tools to get rid of bone spurs, remove loose bodies, or smooth the cartilage. Your surgeon may also do lavage in the joint. Lavage involves rinsing the joint with a sterile saltwater solution. Lavage helps remove tiny debris that may be irritating the joint. The arthroscope allows the surgeon to watch what he or she is doing on a TV screen during the procedure.

The majority of patients treated with arthroscopic debridement for elbow osteoarthritis have less pain and more motion after surgery. Symptoms may come back in some patients, but they are usually less severe.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS


SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

Interposition Arthroplasty

Before the invention of high-quality artificial joints, surgeons used many techniques to keep the bone surfaces of arthritic joints from rubbing against each other. One of these techniques is interposition arthroplasty.




This procedure involves placing a piece of tendon or fascia between the bony surface of the elbow joint. (Fascia is a connective tissue, similar to tendon, that lies in a flat sheet. It covers the muscles and acts as a divider between different compartments of the body.)

As the joint heals, the tendon or fascia forms a cushion of thick, tough tissue between the bones. The tissue pads the ends of the bones and reduces pain while still allowing the elbow to move.

Interposition arthroplasty is still a useful procedure in some cases. It works fairly well in the elbow. It doesn't work very well in the weight-bearing joints of the hip, knee, and ankle.

Elbow Fusion


A fusion surgery (also called arthrodesis) eliminates pain by making the bones of the joint grow together, or fuse, into one solid bone. Fusions were very common before the invention of artificial joints. Even today, joint fusions are commonly used



in many different joints to get rid of the pain of arthritis. An elbow fusion will greatly decrease the motion in your arm. However, it does leave you with a strong and pain-free elbow. People who need a good range of motion in their elbow should consider another type of operation, such as an elbow joint replacement.

Elbow Joint Replacement

Elbow joint replacement is not nearly as common as hip, knee, or shoulder replacement. This is true for a couple of reasons. Osteoarthritis in the elbow is not as common as osteoarthritis in weight-bearing joints.



NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

Elbow joint replacement also has a higher complication rate than the more common replacement surgeries. Infection and slowed healing in the surgical incision are two complications of this type of procedure.

The elbow joint replacement is a good choice for patients who need improved motion rather than strength. Older patients who don't need as much strength will probably prefer the results of elbow replacement surgery. Patients with advanced rheumatoid arthritis are also good candidates for elbow joint replacement.

REHABILITATION

When will I be able to use my elbow again?

Non-Surgical Rehabilitation

If you don't need surgery, range-of-motion exercises for the elbow should be started as

pain eases. These exercises are followed by a program of strengthening that may include shoulder and upper back exercises. You'll be given tips on keeping your symptoms under control. You will probably progress to a home program within four to six weeks.

AFTER SURGERY

Your elbow will be bandaged with a well-padded dressing and an elbow splint for support. Physical or occupational therapy sessions may be needed for up to three months after surgery. The first few treatment sessions will focus on controlling the pain and swelling from surgery.

You will then begin to do exercises that help strengthen and stabilise the muscles around the joint. Your therapist will give you tips on ways to do your activities without straining your elbow.

NEXT PAGE ►

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com

NOTES:

ABOUT MR GEORGE-MALAL

Mr Joby George-Malal is an experienced Consultant Orthopaedic Surgeon with special interest, training and expertise in upper limb (shoulder/ elbow/ hand) problems.

CLINICS

SPIRE HARPENDEN HOSPITAL, HARPENDEN
THE SAXON CLINIC, MILTON KEYNES
THE MANOR HOSPITAL, BEDFORD

CONTACT

EMAIL: info@georgemalal.com
TEL: 03330 124 051
WEB: georgemalal.com